

MYCOTAXON

<http://dx.doi.org/10.5248/117.423>

Volume 117, pp. 423–427

July–September 2011

First record of *Resupinatus poriaeformis* (Agaricomycetes) from South AmericaGEORGEA SANTOS NOGUEIRA-MELO^{1*}, LEIF RYVARDEN²
& TATIANA BAPTISTA GIBERTONI¹¹ Universidade Federal de Pernambuco, Departamento de Micologia,
Av. Nelson Chaves s/n, CEP 50760-420, Recife, PE, Brazil² University of Oslo, Department of Botany, P. O. Box 1045, Blindern, N-0316, Oslo, Norway*CORRESPONDENCE TO: georgeacomea@hotmail.com

ABSTRACT — *Resupinatus poriaeformis* is reported as new to South America. Description of the species and a key to the accepted species of the genus reported to South America are provided.

KEY WORDS — *Tricholomataceae*, diversity

Introduction

Resupinatus was validated by Gray in 1821 and comprises about 15–20 valid species (www.indexfungorum.org; <http://www.cbs.knaw.nl>; Kirk et al. 2008). It is a cosmopolitan genus with small, resupinate, or pendent basidiomata, a partially gelatinized context, and clamped hyphae (Singer 1962; Thorn 1986; Pegler 1986). Recently Thorn et al. (2005) placed some cyphelloid fungi or “reduced agarics” with cup-shaped basidiomata into *Resupinatus* based on molecular and morphological studies.

Although some species of *Agaricales* have been registered in mangroves, there are no reports of *Resupinatus* from this ecosystem (Hibbett & Binder 2001; Baltazar et al. 2009). However there are five reports of the genus from South America: *R. alboniger* (Pat.) Singer 1978, *R. applicatus* (Batsch) Gray 1821, *R. dealbatus* (Berk.) Singer 1973, *R. graminum* (Singer) Singer 1973, and *R. hyalinus* (Singer) Thorn et al. 2005. Of these, *R. applicatus*, *R. dealbatus*, and *R. hyalinus* occur in Brazil (Singer 1943; 1973; 1989; Pegler 1997; Thorn & Barron 1986). We present a description of a new occurrence and a key to the *Resupinatus* species in South America.

Material & methods

Brazilian mangroves extend from 4°30'N to 28°30'S and despite the limited floristic diversity vary significantly in plant growth form, species distribution patterns, and stand structure (Schaeffer-Novelli et al. 1990). From March 2009 to March 2010, 36 field trips were undertaken in four mangroves in Pernambuco state in northeast Brazil: Maria Farinha (07°51'24.8"S 34°50'32.7"W), Itamaracá (07°46'52.6"S 34°52'53.3"W), Maracaípe (08°32'22.8"S 35°00'29.1"W), and Rio Formoso (08°41'20.8"S 35°06'06.6"W).

After collection, the specimen was analyzed macro- (shape, color, hymenial surface) and micromorphologically (hyphal system, presence/absence and measurements of sterile structures and basidiospores). Microscopical observations were made from slide preparations with 5% KOH, stained with 1% of aqueous phloxine, and Melzer's reagent (Ryvardeen 1991). Color designation followed Watling (1969). The material was incorporated to URM.

Scanning electron microscopy (SEM) was conducted at the Laboratório de Microscopia Eletrônica (DF/UFPE). Sections were removed from dried basidiomata and mounted directly on aluminum stubs using carbon adhesive tabs. The fragments were coated with 8–13 nm of gold using a Baltec SCD050 sputter coater and examined with a JEOL JSM-5900 scanning electron microscope.

Taxonomy

Resupinatus poriaeformis (Pers.) Thorn, Moncalvo & Redhead,

Mycologia 97(5): 1148 (2006)

FIGURE 1–2

= *Peziza anomala* var. *poriaeformis* Pers., Syn. meth. fung. 2: 656 (1801)

BASIDIOMATA annual, resupinate, cup-shaped, up to 2–3mm, gregarious, subiculum tomentose to cottony, up to 0.1 mm, grayish (smoke grey 34 to mouse grey 35), without reaction in KOH. **HYMENIAL SURFACE** smooth, dark grey (drab 33 to mouse grey 35). **HYPHAL SYSTEM** monomitic; generative hyphae hyaline, clamped, thin-walled. **BASIDIA** hyaline, clavate, guttulate, 25–30 × 5–7 µm, 4-sterigmata. **CYSTIDIA** absent. **DENDROHYPHIDIA** present. **BASIDIOSPORES** globose, hyaline, thin-walled, smooth, inamyloid, acyanophilous, 4.5–5 µm in diam.

MATERIAL EXAMINED: BRAZIL. PERNAMBUCO: Mangrove of Maria Farinha, on dead *Rhizophora mangle* L. (*Rhizophoraceae*), 28.VI.2009, G.S. Nogueira-Melo et al. NM006 (URM 82257).

REMARKS: *Resupinatus poriaeformis* is characterized by small grey cup-shaped basidiomata and globose basidiospores (4.5–5 µm diam., subglobose and ≤ 6 µm in the literature). The species resembles *R. hyalinus* based on the basidioma size and smooth hymenial surface. However, the basidiospores of *R. hyalinus* are elliptical (6–6.5 × 3–3.4 µm). The other species reported from South America have larger basidiomata (0.5–4 cm) and lamellate hymenial surface. Additionally, the basidiospores are ellipsoid (5.8–7.8 × 2.7–3.6 µm) in *R. alboniger*, cylindrical (5–8 × 2.5–3 µm) in *R. dealbatus*, and subglobose (4.4–6




FIG. 1–2. *Resupinatus poriaeformis*. 1. Basidiospores, 2. Basidiomata. Scale bar = 2 mm.

× 4–4.8 µm) in *R. applicatus* (Corner 1981; Ryvar den & Gilbertson 1993; Singer 1943, 1962; Thorn et al. 1986).

DISTRIBUTION — Ryvar den & Gilbertson (1993) and Bernicchia (2005) report this species as cosmopolitan. Records are known from Honduras, Italy, New Zealand, Puerto Rico, South Africa and Sweden (Cunningham 1963;

Ryvar den 1988; Ryvar den & Gilbertson 1993; Pennycook & Galloway 2004; Thorn et al. 1986, 2006; Bernicchia 2005, O database).

Key to the *Resupinatus* species recorded from South America

- 1a. Hymenial surface lamellate 2
- 1b. Hymenial surface smooth 5
- 2a. Basidiomata reniform to flabelliform up to 2.5–4 cm *R. dealbatus*
- 2b. Basidiomata cupulate to flabelliform up to 1 cm 3
- 3a. Cystidial elements lacking *R. graminum*
- 3b. Cystidial elements present 4
- 4a. Basidiospores subglobose, 4–4.5 x 5 µm *R. applicatus*
- 4a. Basidiospores ellipsoid to cylindrical, slightly curved 5.8–7.8 x 2.7–3.6 µm
..... *R. alboniger*
- 5a. Basidiospores globose to subglobose, 4.5 x 6 µm diam. *R. poriaeformis*
- 5b. Basidiospores ellipsoid, 6–6.5 x 3.3–4 µm *R. hyalinus*

Acknowledgments

We would like to thank Dr. A. Bernicchia and Dr. R.G. Thorn for critically reviewing the manuscript, Sérgio Santos for assistance with the SEM, and Leandro Agra for taking the photo. Further, we acknowledge the CNPq for the master scholarship of GSNM, the FACEPE (APQ 0444-2.03/08) and the Pós-Graduação em Biologia de Fungos (UFPE) for financial support.

Literature cited

Baltazar JM, Trierweiler-Pereira L, Loguercio-Leite C. 2009. A checklist of xylophilous basidiomycetes (*Basidiomycota*) in mangroves. *Mycotaxon* 107: 221–224.

Bernicchia A. 2005. *Polyporaceae* s.l. Edizione Candusso, Alassio. 808 p.

Corner E.J.H. 1981. The agaric genus *Lentinus*, *Panus* and *Pleurotus* with particular reference to Malaysian species. J. Cramer, Germany. 169 p.

Cunningham G.H. 1963. The *Thelephoraceae* of Australia and New Zealand. New Zealand Department of Scientific and Industrial Research, Bulletin 145. 359 p.

Hibbett DS, Binder M. 2001. Evolution of marine mushrooms. *Biological Bulletin* 201: 319–322.

Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. Dictionary of the fungi. 10th ed. CABI Publishing, Wallingford. 771 p.

Pegler DN. 1986. Agaric flora of Sri Lanka. *Kew Bulletin Additional Series* 12. 519 p.

Pegler DN. 1997. The agarics of São Paulo, Brazil. Royal Botanic Garden, Kew. 68 p.

Pennycook SR, Galloway DJ. 2004. Checklist of New Zealand “fungi”: 401–488, in: Introduction to fungi of New Zealand. Fungi of New Zealand/Ngā Harore o Aotearoa, vol. 1. Fungal Diversity Research Series 14.

Ryvar den L. 1988. Type studies in the *Polyporaceae* 19. Species described by M.C. Cooke. *Mycotaxon* 31: 45–58.

Ryvar den L. 1991. Genera of polypores – nomenclature and taxonomy. *Synopsis Fungorum* 5: 1–363.

Ryvar den L, Gilbertson RL. 1993. European polypores, vol. 2. Fungiflora, Oslo. pp. 394–743.

- Schaeffer-Novelli Y, Citrón-Molero G, Adaime RR. 1990. Variability of mangrove ecosystems along the Brazilian coast. *Estuaries* 13(2): 204–218.
- Singer R. 1943. Type studies on *Basidiomycetes* II. *Mycologia* 32(2): 142–163.
- Singer R. 1962. The *Agaricales* in modern taxonomy. 2nd ed. Germany J: Cramer. 915 p.
- Singer R. 1973. Diagnoses fungorum novorum agaricalium III. Beihefte zur Sydowia 7. 106 p.
- Singer R. 1989. New taxa and new combinations of *Agaricales* (Diagnoses fungorum novorum agaricalium IV). *Fieldiana Botany* 21: 1–133.
- Thorn RG, Barron GL. 1986. *Nematoctonus* and the tribe *Resupinateae* in Ontario, Canada. *Mycotaxon* 25(2): 321–453.
- Thorn RG, Moncalvo JM, Redhead SA, Lodge DJ, Martín MP. 2005. A new poroid species of *Resupinatus* from Puerto Rico, with a reassessment of the cyphelloid genus *Stigmatolemma*. *Mycologia* 97(5): 1140–1151. <http://dx.doi.org/10.3852/mycologia.97.5.1140>
- Watling R. 1969. Colour identification chart. Her Majesty's Stationary Office, Edinburgh.